



Comparative Analysis of Antibacterial Effect of Ethanol Extracted *Allium sativum* (Garlic) Oil and Ampicillin on *Bacillus cereus*, *Bacillus subtilis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Escherichia coli*

¹ADAMU, Jacob Eshiemokhai, ¹OGUNSUMI, ¹Akintunde Isreal; ¹IGBOAMA, Magdalene C, ¹SULEIMAN, Ismail O and ²ILESANMI, Idayat Adedayo

¹Department of Biological Sciences, Federal Polytechnic Ede, Osun State

²Department of Environmental Health, University of Ibadan, Oyo State

Corresponding Author Email Address & Contact: jacobeshiemokhai@gmail.com; +2347039393768

Abstract: Antimicrobial effect of the oil extracted from the bulbs of *Allium sativum* (garlic) purchased from the “Oje” market of Edeland in Osun State, Nigeria was assessed. The oil was extracted by Soxhlet extraction method using ethanol as extraction solvent. The extract was screened against five clinical microorganisms; *Escherichia coli*, *Bacillus cereus*, *Bacillus subtilis*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* in-vitro, using agar-well-diffusion method and Ampicillin as control. The extracted oil is yellowish in colour with a pungent smell. Using different dilution concentrations of the extract, the highest zone of inhibition was recorded against *B. subtilis* 27±1.2 while *P. aeruginosa* showed least sensitivity at 8.0±0.0 at 100%. At 75% dilution, the highest zone of inhibition was 20±1.2 against *E. coli* while the least was against *P. aeruginosa* 13±1.4. At 25%, highest zone of inhibition was 18±0.0 against *B. subtilis* and least 11±1.2. The effect of ampicillin ranges between 24±1.4 and 16±1.4 against the tested microorganisms. The ethanol extract of *Allium sativum* showed more sensitivity to *B. subtilis*, at 100%, and *E. coli*, at 100%, 75%, and 25%, compared to Ampicillin.

Keywords: Antibacterial, Ampicillin, Extract, Garlic.

INTRODUCTION

The use of medicinal plants and its products to treat infections is an age-long practice, in Africa. Medicinal plants are important part of our natural wealth where they serve as important therapeutic agents as well as valuable raw materials for manufacturing numerous traditional and conventional medicines (krishnaiah *et al.*, 2009). Many indigenous plants in Nigeria are used in inhibiting and curing various life threatening infections or diseases caused by pathogenic microorganisms. Medicinal plants are invariably used to source for new drugs, many drugs which are in use today were either obtained from plants or developed using plants chemical structure as templates (Ajaiyeoba *et al.*, 2006). The importance of locally available plants as medicine for treatment of infectious diseases cannot be underestimated (Oladeji, 2016). It is therefore significantly important to examine the phyto-chemical properties and antimicrobial effects of some medicinal plants commonly used in Nigeria.

Interest in plants with antimicrobial properties is being reviewed because of the emergence of resistance strains of disease causing microorganisms against some conventional antimicrobials drugs, (Hawassa & Sekota, 2016). The World Health Organization (WHO, 2001) estimated that the world population that uses medicinal plants to treat different forms of ailments and diseases is about 80% and this value is even much higher in African. Some plant produce essential oils, valuable chemical compound which are considered the chemical weapons. These compounds may dissuade insects or protect the plant against bacterial and fungal attacks while they as well act as instrument to attract and seduce their pollinators (Hamid *et al.*, 2011). He further stated that these volatile liquid's aromatic compounds may be found in different parts hitherto in the leaves (oregano), seed (almond), flower (jasmine), peel (bergamot), berries (juniper), rhizome (galangal ginger), root (angelica archangelica), bark (sassafras), wood (agar wood), resin (frankincense), petals (rose), and bulb (garlic).

Garlic (*Allium sativum*) has had an important dietary and medicinal role for centuries. It is a large annual plant of the Liliaceae family, which grows in most of African countries where it is used in traditional medicine for

infectious disease and some other cases (Daka, 2011). It is an annual plant with superficial adventitious roots, bulbs composed of a disk like stem (Sofoworo, A., 1982). Garlic played important dietary and medicinal role for centuries where it is commonly used in food as flavoring agent and when consumed in food it does not produce any obvious side effects (Padiya and Banergee, 2013). It was reported that *Allium sativum* as a medicinal plant portray some anti bacterial effects against some organisms but it is under-utilized due to its odour and taste, although, garlic is still being employed in folk medicine all over the world for the treatment of a variety of diseases (Ali, Thompson and Afzal, 2000). The prophylactic and therapeutic effects of garlic are ascribed to specific oil and water soluble organo-sulfur compounds, which are responsible for its typical odor and flavor (Mims et al., 2021). Venegas et al. (2016) also reported the garlic bacterial inhibitory effect on species of *Aerobacter*, *Aeromonas*, *Bacillus*, *Citrobacter*, *Clostridium*, *Enterobacter*, *Escherichia*, *Klebsiella*, *Lactobacillus*, *Leuconostoc*, *Micrococcus*, *Mycobacterium*, *Proteus*, *Pseudomonas*, *Salmonella*, *Shigella*, *Staphylococcus*, *Streptococcus* and *Vibrio*.

Garlic act as an anti-bacterial agent that inhibits the growth of infectious agents while at the same time protect the body from pathogens, an example is the deadly *Bacillus anthracis* which causes the disease anthrax (Wolde et al, 2018). Dini, Fabbri and Geraci (2011) *Mycobacterium tuberculosis*, a pathogen that causes tuberculosis showed successfully sensitivity to several sulphur components found in Garlic. More so, Garlic consumption was reported to have significant effects on lowering blood pressure, prevention of atherosclerosis, reduction of serum cholesterol and triglyceride, inhibition of platelet aggregation, and increasing fibrinolytic activity (Kumar et al., 2013). Furthermore, Leyla, Peir and Ali, (2014) stated that in *in-vivo* animal experiments, intravenous administration of garlic extracts produced slight reduction in both systolic and diastolic pressures and oral ingestion of garlic extract in hypertensive animals brought the blood pressure back to the normal level (Marmitt, Goettert and Rempel, 2021).). In addition, the preventive effect of garlic on atherosclerosis has been attributed to its capacity to reduce lipid content in arterial membrane (Yu-Yah and Liu, 2001). It was also reported that garlic decreases the risk of peripheral arterial occlusive diseases, plasma viscosity, and unstable angina and increases elastic property of blood vessels and capillary perfusion (Allison et al., 2012). More recently, garlic has been proven to be effective against a plethora of gram-positive, gram-negative, and acid-fast bacteria. These include *Salmonella* and *Escherichia coli* (Adler and Beuchat, 2002). O’Gara et al. (2000) documented that garlic shows differential inhibition in beneficial intestinal micro-flora and potentially harmful entero-bacteria which includes *Pseudomonas*, *Proteus*, *Staphylococcus aureus*, *Escherichia coli*, *Salmonella*, *Klebsiella spp.*, *Micrococcus spp.*, *Bacillus subtilis*, *Clostridium spp.*, *Mycobacterium spp.*, and *Helicobacter spp.*

The exploitation and general acceptance of plants of medicinal importance, like *Allium sativa*, is poorly inclined among the elite and the continuous use of chemical based orthodox medicine to fight against pathogenic bacteria has proven to have short or long term effects with the emergence of antibiotics resistant strains which is evident globally. These necessitated the need for alternative antibiotics that is new, natural, plant based, non-toxic, cost effective with minimal or no side effects to eliminate the menace of orthodox medicine. This research focused on the extraction of oil from the bulb of natural garlic plant using ethanol solvents and accessing the antimicrobial effects of the extract on selected pathogenic microorganisms namely *Bacillus subtilis*, *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*.

The aim of this study is to affirm the antimicrobial sensitivity of garlic oil extract on some clinical microorganisms: *Escherichia. coli*, *Bacillus cereus*, *Bacillus Subtilis*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* while the specific objectives assessed physico-chemical component of *Allium sativa* (garlic) oil; assessed the antibacterial effect of garlic oil extract on the named microorganisms; and, compare the sensitivity effect of garlic oil extract with a known antimicrobial agent (Ampicilin).

MATERIAL AND METHODS

Collection of plant material and oil Extraction

Fresh samples of garlic bulbs (*Allium sativum*) were purchased from ‘oje’ market in Ede, Osun state. Aseptically, the cloves on the bulbs were peeled to obtain the edible portion which was diced into tiny bits, dried, ground to powder and stored for use appropriately. In extraction, 15g each of dried garlic powder was weighed and wrapped in a whatman filter paper. Two of the wrapped samples were placed in the thimble. 200ml of ethanol was transferred into the round bottom flask. The soxhlet apparatus was placed on the heating mantle and the ethanol was allowed to boil. The extraction process continued into condensation and the condensing unit was removed from the extraction unit. The oil was obtained after the solvent was evaporated by placing over a

water bath for about 2-3 hours under reduced temperature (50°C) and refluxing at 70°C to remove excess solvent.

Physico-chemical analysis of garlic oil extract

Saponification value of the garlic oil was estimated by the ASTM D464 titration method as described by Abdulhamid *et al.*, (2014). The refractive index of the crude oils samples were measured using an Abbe Refractometer at 30 ± 0.1 after calibration. The iodine value of the oils were determined using a mathematical relationship between refractive index and estimated iodine as being described by Perkins (1995) and reported by Amos *et al.*, (2013). And, the peroxide values of the extracted oils were determined using the standard method described by Akpan *et al.*, (2006).

Preparation of oil concentrates

The extracted oil (100% concentrate) was diluted into different concentration of 75%, 50% and 25%. The dilution was done with methanol. The 100% oil constituted the undiluted form of the oil. The 75% concentration of the oil extract was constituted using 0.75ml of the extracted oil and 0.25ml of methanol in a sterile test tube. The 50% concentration of the oil was constituted using 0.5ml of extracted oil and 0.5ml of methanol in a sterile test tube while the 25% oil concentrate was constituted using 0.25ml of the extracted oil and 0.75ml of methanol in a sterile test tube. The extracted oil at different concentration was stored in a refrigerator at 40°C for further use.

Test organisms

Test microorganisms used were sub-cultured from clinically important microorganisms collected from the Federal Institute of Industrial Research, Oshodi (FIIRO), Lagos Nigeria. The organisms used were *Bacillus cereus* ATCC 10872, *Bacillus subtilis* ATCC 6633, *Staphylococcus aureus* ATCC 25923, *Pseudomonas aeruginosa* ATCC 9027, and *Escherichia coli* ATCC 25922.

Preparation of media

Mueller Hilton agar (MHA) was used for the antimicrobial assay using agar well diffusion method while Nutrient broth was used for sub-culturing test organisms. The media were prepared according to manufacturer's specification. The glass wares used for media preparation were properly washed and rinsed with distilled water and dried-sterilized in the hot air oven at 160°C. Media were sterilized by autoclaving at 121°C for 15 minutes.

Antimicrobial susceptibility testing

The screening was done using agar well diffusion method as described by Harold (2002). Plate culture of 18-hour old broth culture of test organisms was made using pour plate method on a well labeled Mueller Hilton Agar plates using sterile swab and duplicated appropriately in an aseptic manner. Holes were made on the solidified nutrient plates containing the test organisms using a 6mm cork borer. The essential oil in different concentration/dilution was dispensed into the bored holes (100%, 75% and 50% concentrations) while negative control (methanol) and positive control (Ampicillin) dispensed into their appropriate holes. The inoculated plates were incubated at 37°C for 24hours. The resulted clear zones inhibition in millimeters were measured, recorded and inhibitive effects of garlic oil extract estimated.

RESULTS AND DISCUSSION

The physico-chemical features of the extracted oil is presented in Table 1 while Table 2 shows the antimicrobial activities of garlic oil extract against different strains of clinical bacterial.

Table 1: Physico-chemical properties of ethanol extract of *Allium sativa* (garlic) oil.

Parameters	Concentration unit
Specific gravity	0.89 g/ cm ³
Moisture	8.95%
Saponification value	158.1 mg (KOH)/g
Iodine value	11.9 mg/100g
Free fatty acids	1.4±0.1%
Acid value	13.8 mg(KOH)/g

Peroxide value	215 mg/100g
Refractive index	0.97
Melting point	-12°C
Colour	Light yellow
Odour	Pungent

Specific gravity of garlic oil was found to be 0.894g/ cm³ which was lower than most edible oil such as sunflower oil, cotton seed oil and olive oil reported by Rahman and Lowe (2006) to be approximately 0.919g/ cm³. The other features of the extracted oil was found to include; 8.95% moisture, 0.97 refractive index, 11.9 iodine level, acid value (13.8), free fatty acids (1.4%), saponification (158.1) and peroxide values (215) respectively.

The oil from *Allium sativum* exhibited high activity against most of the organisms. The largest zone of inhibition was 27 ± 1.21 recorded against *Bacillus subtilis* ATCC 6633 while the least zone of inhibition was 8 ± 0.00 recorded against *Pseudomonas aeruginosa* ATCC 9027 at 100% concentration. The activity of the oil at 75% concentration was significant against the bacteria species; *Escherichia coli* ATCC 25922 was highly inhibited at 20 ± 1.2, followed by *Bacillus cereus* ATCC 6633 at 18 ± 1.06 while 15 ± 1.05, 13 ± 1.41 and 15 ± 4.24 was recorded against *Bacillus subtilis* ATCC 10872, *Pseudomonas aeruginosa* ATCC9027 and *Staphylococcus aureus* ATCC25922 respectively. At 25% concentration, zones of inhibition of 18 ± 0.00, 12 ± 1.41; 13 ± 0.00 11 ± 1.2 and 14 ± 0.00 was recorded against *Bacillus subtilis* ATCC 6633, *Bacillus cereus* ATCC 10872, *Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 9027 and *Staphylococcus aureus* ATCC 25923 respectively.

Table 2: Antimicrobial effect of ethanol extract of *Allium sativa* (garlic) oil

Microorganism	Z o n e o f i n h i b i t i o n (m m)			
	Essential oil 1 0 0 %	Essential oil 7 5 %	Essential oil 2 5 %	A m p i c i l l i n
<i>Bacillus subtilis</i> A T C C 6 6 3 3	2 7 ± 1 . 2 1	1 5 ± 1 . 0 5	1 8 ± 0 . 0 0	2 0 ± 0 . 0 0
<i>Bacillus cereus</i> A T C C 1 0 8 7 2	1 3 ± 1 . 0 0	1 8 ± 1 . 0 8	1 2 ± 1 . 4 1	2 2 ± 4 . 2 4
<i>Escherichia coli</i> A T C C 2 5 9 2 2	2 1 ± 1 . 5 0	2 0 ± 1 . 2 0	1 3 ± 0 . 0 0	1 6 ± 1 . 4 1
<i>P. aeruginosa</i> A T C C 9 0 2 7	8 ± 0 . 0 0	1 3 ± 1 . 4 1	1 1 ± 1 . 2 0	1 6 ± 2 . 8 3
<i>S. aureus</i> A T C C 2 5 9 2 2	2 2 ± 2 . 8 3	1 5 ± 4 . 2 4	1 4 ± 0 . 0 0	2 4 ± 1 . 4 1

The garlic oil at 100%, 75% and 25% showed good activity against bacteria species when compared with the standard antibiotic ampicillin.

The result of the antimicrobial effect of the ethanolic extract of *Allium sativum* revealed the largest zone of inhibition of 27 ± 1.21 against *Bacillus subtilis* ATCC 6633 at 100% concentration which was better than the effect of ampicillin against the microorganism. Also, the oil extract was more sensitive to *Escherichia coli* ATCC 25922 compared to Ampicillin. Daka (2011) reported that extract of garlic showed antibacterial activity against *Staphylococcus aureus in vitro*. In addition, Manu, Rachna and Nidhi (2019) study also suggested that garlic have anti -bacterial activity against *Staphylococcus aureus* and *Escherichia coli* and can be used against pathogenic microorganisms.




Figure 1: Comparative chart showing Antimicrobial effects of ethanol extract *Allium sativa* (garlic) oil and Ampicillin

The oil extract was more sensitive against *E. coli* at all concentration compared to Ampicillin while *B. subtilis* showed better susceptibility at 100% concentration compared to Ampicillin. Whereas, other organisms *B. cereus*, *P. aeruginosa* and *S. aureus* were less susceptible to the oil extract at all concentration compared to Ampicillin. Onyeagba reported a contrary result that crude extracts of garlic applied singly did not exhibit any *in vitro* inhibition on the growth of test organisms including *Staphylococcus* spp. (Onyeagba, 2004). But, Hamdy, Soad and Doaa (2016), reported that garlic essential oil extracts exhibited different inhibition levels against different tested bacteria. Also, Daka (2011) reported that oil extract of garlic have concentration dependent antibacterial effect against *Staphylococcus aureus*.

The result showed that extracted oil from the bulbs of *Allium sativum* (garlic) exhibit significant activity against tested microorganisms *Bacillus subtilis*, *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*, although, with different level of severity. The inhibitory effect of the essential oil at different percentages (100%, 75%, and 25%) showed relatively high zones against *B. subtilis*, *B. cereus*, *E. coli* and *S. aureus* except for *Pseudomonas aeruginosa* whose zone of inhibition was comparably very weak compared to the standard Ampicillin. In the report by Hamdy, Soad and Doaa (2016) it was also affirmed that garlic oil extract exhibit variant level of antimicrobial activity against different species of microorganism and it further declared that garlic oil extracts exhibited marked inhibition activity against bacteria at high concentrations. The findings suggested that the garlic oil extracted with ethanol solvents showed appreciable antimicrobial activities against *Bacillus cereus*, *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. The ethanolic oil extracts had more inhibitory effect on *E. coli* at 100%, 75% and 25% dilution while *B. subtilis* was more susceptible to the *allium sativum* oil at 100% dilution compared to Ampicillin. Study by Elias *et.al.* (2022) showed that significance differences were recorded in the inhibitory effect of garlic oil extracts at different concentration levels against different bacteria.

CONCLUSION AND RECOMMENDATION

In conclusion, ethanolic extract garlic oil have antibacterial potential and could be used as an alternative to common antibacterial drugs in the treatment of some common bacterial infections. Albeit, expanded research into this subject could lead to more discovery. It is recommended to carry out further trials to establish the clinical efficacy, safety, antioxidant effect and toxicity of the extracted garlic oil.

REFERENCES

- Ademiluyi, A. O., Oboh, G., Owoloye, T. R., & Agbebi, O. J. (2013). Modulatory effects of dietary inclusion of garlic (*Allium sativum*) on gentamycin-induced hepatotoxicity and oxidative stress in rats. *Asian Pacific journal of tropical biomedicine*, 3(6), 470-475. [https://doi.org/10.1016/S2221-1691\(13\)60098-2](https://doi.org/10.1016/S2221-1691(13)60098-2)
- Adler, B. B., & Beuchat, L. R. (2002). Death of *Salmonella*, *Escherichia coli* O157: H7, and *Listeria monocytogenes* in garlic butter as affected by storage temperature. *Journal of food protection*, 65(12), 1976-1980. <https://doi.org/10.4315/0362-028X-65.12.1976>
- Ajaiyeoba, E., Falade, M., Ogbale, O., Okpako, L., & Akinboye, D. (2006). In vivo antimalarial and cytotoxic properties of *Annona senegalensis* extract. *African Journal of Traditional, Complementary and Alternative Medicines*, 3(1), 137-141.
- Ali, M., Thomson, M., & Afzal, M. (2000). Garlic and onions: their effect on eicosanoid metabolism and its clinical relevance. *Prostaglandins, Leukotrienes and Essential Fatty Acids (PLEFA)*, 62(2), 55-73. <https://doi.org/10.1054/plf.1999.0124>
- Allison, G. L., Lowe, G. M., & Rahman, K. (2012). Aged garlic extract inhibits platelet activation by increasing intracellular cAMP and reducing the interaction of GPIIb / IIIa receptor with fibrinogen. *Lifesciences*, 91(25-26), 1275-1280. <https://doi.org/10.1016/j.lfs.2012.09.019>
- Ashraf, R., Aamir, K., Shaikh, A. R., & Ahmed, T. (2005). Effects of garlic on dyslipidemia in patients with type 2 diabetes mellitus. *J Ayub Med Coll Abbottabad*, 17(3), 60-64.
- Atlas R. M., (1997). Principles of microbiology. McGraw Hill, New York, USA.
- Bakhshi, M., Taheri, J. B., Basir Shabestari, S., Tanik, A., & Pahlevan, R. (2012). Comparison of therapeutic effect of aqueous extract of garlic and nystatin mouthwash in denture stomatitis. *Gerodontology*, 29(2), e680-e684. <https://doi.org/10.1111/j.1741-2358.2011.00544>
- Banerjee, S. K., & Maulik, S. K. (2002). Effect of garlic on cardiovascular disorders: a review. *Nutrition journal*, 1(1), 1-14. <https://doi.org/10.1186/1475-2891-1-4>.
- Capasso, A. (2013). Antioxidant action and therapeutic efficacy of *Allium sativum* L. *Molecules*, 18(1), 690-700. <https://doi.org/10.3390/molecules18010690>
- Daka Deresse, (2011): Antibacterial effect of garlic (*Allium sativum*) on *Staphylococcus aureus*: An in vitro study: African Journal of Biotechnology Vol. 10 (4), pp. 666-669, 24 January, 2011 Available online at <http://www.academicjournals.org/AJB> DOI: 10.5897/AJB09.553 ISSN 1684-5315 © 2011 Academic Journals
- Dini, C., Fabbri, A., & Geraci, A. (2011). The potential role of garlic (*Allium sativum*) against the multi-drug resistant tuberculosis pandemic: a review. *Annali dell'Istituto superiore di sanità*, 47, 465-473.
- Elias L., Zekeria Y., Mulugeta D., Sultan S., Megersa I., Sewnet M., and Jemal T. (2022): Physicochemical Properties and Biological Activities of Garlic (*Allium sativum* L.) Bulb and Leek (*Allium ampeloprasum* L) Leaf Oil Extracts. The Scientific World Journal; Volume 2022, Article ID 6573754, 7 pages; <https://doi.org/10.1155/2022/6573754>.
- Hamdy R Soltan, Soad M Ahmed, Doaa A Emam (2016): Comparative antibacterial activity of garlic essential oil extracted by hydro-distillation and diethyl ether extraction methods on four pathogenic bacteria. MedCare Journal of Advances in Plants & Agriculture Research, Volume 4 Issue 2, Published: July 21, 2016.

- Hamid, A. A., Aiyelaagbe, O. O., and Usman, L. A. (2011). Essential oils: its medicinal and pharmacological uses. *International Journal of Current Research*, 33(2): 86 – 89.
- Harold, S. B. (2002). *Laboratory manual in General microbiology*. McGraw Hill, New York, USA.
- Hawassa, E., & Sekota, E. (2016). Anti-Bacterial Effect of Garlic (*Allium sativum*) against Clinical Isolates of *Staphylococcus aureus* and *Escherichia coli* from Patients Attending Hawassa Referral Hospital, Ethiopia. *Journal of Infectious Diseases*, 2(2), 18. DOI: 10.21767/2472-1093.100023
- Houshmand, B., Mahjour, F., Dianat, O. (2013). Antibacterial effect of different concentrations of garlic (*Allium sativum*) extract on dental plaque bacteria. *Indian Journal Dental Research*. 24:71
- Jonkers, D., Sluimer, J., Stobberingh, E. (1999). Effect of garlic on vancomycin-resistant enterococci. *Antimicrobial Agents and Chemotherapy*. 43:3045
- Krishnaiah, D., Devi, T., Bomo, A. and Sarbatly, R. (2009). Studies on Phytochemical constituents of Six Malaysian Medicinal Plants. *Journal of Medicinal plants Research*. 3(2); 67-72.
- Kumar, R, Chandra, S., Samsher, Kumar, K., Kumar, T., Kumar V. (2016). Analysis of the physical and chemical characteristics of edible blended oils. *United states department of Agriculture*. 2:385-393.
- Kumar, R., Chhatwal, S., Arora, S., Sharma, S., Singh, J., Singh, N., Bhandari V., Khurana A. (2013). Antihyperglycemic, antihyperlipidemic, anti-inflammatory and adenosine deaminase–lowering effects of garlic in patients with type 2 diabetes mellitus with obesity. *Diabetes Metabolic Syndrome and Obesity*. (6):49-56 DOI:10.1016 /j.phytochem.2012.03.009
- Lanzotti, V., Barile, E., Antignani, V., Bonanomi, G., Scala, F. (2012). Antifungal saponins from bulbs of garlic, *Allium sativum* L var. Voghiera. *Phytochemistry*. 78:126-134
- Leyla, B., Peir, H. K., Ali, G., (2014). Garlic: a review of potential therapeutic effects. *Avicenna journal of phytomedicine*.4(11): 1-14
- Lissiman, E., Bhasale, A. L., Cohen, M. (2012). Garlic for the common cold. *Cochrane Database Syst Rev*. (3).
- Manu Yadav, Rachna Bohra, Nidhi Gupta (2019): *In vitro* Determination of Antibacterial Effect of Garlic (*Allium sativum*) on *Staphylococcus aureus* and *E. coli*. *International Journal of Current Microbiology and Applied Sciences* ISSN: 2319-7706 Volume 8 Number 09 (2019).
- Marmitt, D. J., Goettert, M. I., & Rempel, C. (2021). Recent Scientific Production of Garlic (*Allium sativum* L.) confirms important therapeutic properties. *Brazilian Journal of Development*, 7(5), 51030-51055.
- Mims, S., Shreesh, O., Huba, K., Adeghate, E. (2021). Chemical constituents and medicinal properties of *Allium* species. *Journal of Molecular and cellular biochemistry* DOI:10.1007/s11010-021-04213-2
- O'Gara, E. A., Hill, D. J., Maslin, D. J. (2000). Activities of garlic oil, garlic powder, and their diallyl constituents against *Helicobacter pylori*. *Applied Environmental Microbiology*. 66:2269-2273
- Ohaeri, O. C. (2001). Effect of garlic oil on the levels of various enzyme in the serum and tissue of streptozotocin diabetic rats. *Bioscience Reports*. 21:19-24
- Oladeji, O. (2016). The characteristics and roles of medicinal plants: some important medicinal plants in Nigeria. *Natural products Indian journal*. 12(3):102
- Omar, S. H. (2013). Garlic and cardiovascular diseases. In: Ramawat K., (eds) *Natural products*. Springer, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-642-221446-158>.
- Onyeagba R.A., Ugbogu O.C., Okeke C.U., Iroakasi O., (2004): Studies on the antimicrobial effects of garlic (*Allium sativum* Linn), ginger (*Zingiber officinale* Roscoe) and lime (*Citrus aurantifolia* Linn). *Afr. J. Biotechnology* 3 (10): 552-554.

- Padiya, R., Banerjee, S. K. (2013). Garlic as an anti-diabetic agent: recent progress and patent reviews. *Recent Patents on Food, Nutrition & Agriculture*. 5:105-127
- Rahman, K., Lowe, G. M., (2006). Significance of garlic and its constituents in cancer and cardiovascular disease. *The Journal of nutrition*. 136: 736-740.
- Seyed, H.E., Neyed, A. S., Shabani, S. (2014). Antibacterial effect of garlic aqueous extract on staphylococcus aureus in hamburger. *Jundishapur Journal of Microbiology*; 7(11): e13134
- Sofoworo, A. (1982). *Medicinal plants and traditional medicine in Africa*. John Wiley and sons Ltd., New York pp256
- Soltan, H. R., Ahmed, S. M., Emam, D. A. (2016). Comparative antibacterial activity of garlic essential oil extracted by hydrodistillation and diethylether extraction methods on four pathogenic bacteria. *Advances in plants & agriculture research*. 4(2):261-264
- Soxhlet extraction method- estimation of fat in food <http://people.umass.edu>. Cited on 18/08/2021.
- WHO (2001). *Legal status of traditional medicine and complementary/alternative medicine. A worldwide review*. WHO publishing 1.
- Wolde, T., Kuma, H., Trueha, K., Yabeker, A. (2018). Antibacterial activity of garlic extract against human pathogenic bacteria. *Journal of pharmacovigilance* 6(1). DOI: 10.4172/2329-6887.1000253
- Yu-Yan, Y., Liu, L. (2001). Cholesterol lowering effect of garlic extracts and organosulfur compounds: Human and animal studies. *The Journal of Nutrition*. 131: 989-993.
- Venegas, A., Touma, J. H., Bravo, J., & Perez-Perez, G. (2016). Progress in use of natural products and their active components against *Helicobacter pylori*. *Advances in Microbiology*, 6(14), 1091. <https://doi.org/10.4236/aim.2016.614101>